

*Aventis Pasteur*



# Clinical Experience at Aventis Pasteur With Combination Acellular DTP and Hib Conjugate Vaccines

Carlton K. Meschievitz, M.D., M.P.H.  
Vice President, Scientific and Medical Affairs  
Aventis Pasteur

Aventis Pasteur waives confidentiality of the data contained in these presentation materials.  
Aventis Pasteur grants FDA the authority to release these presentation materials to the public  
through electronic and/or other means deemed appropriate.

Carlton Koon Meschievitz MD

2636 '00 JAN 24 P1:56

## Age Distribution and Vaccination Status of Children <5 years of Age with Hib Disease, 1996 and 1997\*

|            | Total Cases | Vaccination Status |            |
|------------|-------------|--------------------|------------|
|            |             | incomplete         | complete   |
| < 6 months | 69 (48%)    | -                  | -          |
| > 6 months | 75 (52%)    | 48 (64%)           | 27 (36%)** |

\*\*primary series = 13 children; primary series + booster = 14 children

\*MMWR, November 27, 1998

*Aventis Pasteur*



## **Alaska Experience\***

Prior to introduction of Hib vaccine (<1991)

- Peak incidence at 4-7 months of age
- 25-40 cases/year
- Carriage (5%)

PedvaxHib® (1991-95)

- Nearly eliminated disease (1-3 cases/year)
- No effect on carriage (8.6%)

Tetramune® (1996-97)

- 10 cases/year

Reasons for reemergence of Hib disease

- PedvaxHib® did not eliminate carriage
- Tetramune® did not protect against early Hib disease

Both the population and the specific Hib vaccine may be important in control of Hib

**\* Galil, et.al. Journal of Infectious Diseases, 1999; 179:101-6.**

## Combination Vaccine Product Identification

TriHIBit<sup>®</sup>

Tripedia<sup>®</sup> (DTaP 2 component) used to reconstitute ActHIB<sup>®</sup>

Quadracel<sup>™</sup>

Tripacel<sup>®</sup> (DTaP 5 component) combined with IPV

Pentacel<sup>™</sup>

Tripacel<sup>®</sup> (DTaP 5 component) combined with IPV used to reconstitute ActHIB<sup>®</sup>

Aventis Pasteur



## TriHIBit<sup>®</sup> in Toddlers\*

|                  | Immunogenicity        |                     |                       |                     |
|------------------|-----------------------|---------------------|-----------------------|---------------------|
|                  | Pre-Dose              |                     | Post-Dose             |                     |
|                  | TriHIBit <sup>®</sup> | Separate Injections | TriHIBit <sup>®</sup> | Separate Injections |
| N                | 88                    | 94                  | 93                    | 98                  |
| Anti-PRP (µg/ml) | 0.89                  | 1.15                | 90.30                 | 80.90               |
| % > 1 µg/ml      | 45.50                 | 53.20               | 100.00                | 100.00              |

\* From FDA approved Product Insert

Anti-PRP Responses in 15 to 20 Month Old Children Following Immunization with TriHIBit<sup>®</sup> Compared to ActHIB<sup>®</sup> and Tripedia<sup>®</sup> Given Concomitantly at Separate Sites\*

*Aventis Pasteur*



## TriHIBit<sup>®</sup> Infants

| Immune Response to PRP Post-Dose 3 |                     |                        |
|------------------------------------|---------------------|------------------------|
| Vaccine                            | Separate            | Combined               |
| N                                  | 69                  | 75                     |
| Anti-PRP                           | 7.0*<br>[5.6 - 8.6] | 4.3<br>[3.0 - 6.0]     |
| % $\geq 0.15$ $\mu\text{g/mL}$     | 100%*               | 94.7%<br>[89.6 - 99.8] |
| % $\geq 1$ $\mu\text{g/mL}$        | 100%†               | 85.3%<br>[77.3 - 93.3] |

\*  $p < 0.05$

†  $p < 0.01$ ; difference of 14.7% (6.7 – 22.7)

Immunogenicity and Lot Consistency of TriHIBit<sup>®</sup> Compared to Tripedia<sup>®</sup> and ActHIB<sup>®</sup> (Study 468-01)

## Adapted from Letter from CBER to Aventis Pasteur

In general, we concur with your definitions for determining the limits for non-inferiority and equivalence for GMT and seroconversion.

Seroconversion - 10% upper limit of the confidence interval

GMT -  $<1.5$  fold difference between combined and separate group

$>2/3$  for ratio of combined to separate groups

*Aventis Pasteur*



## Pentacel™ in Infants Post-Dose 3

| Antigen  | Criteria      | Pentacel™<br>N= 321   | Quadracel™<br>+ActHIB®<br>N= 108 | Comparison<br>Diff. In %<br>(90% CI) |
|----------|---------------|-----------------------|----------------------------------|--------------------------------------|
| Anti-PRP | GMT*          | 4.40<br>(3.78 – 5.13) | 3.83<br>(3.05 - 6.53)            | --                                   |
|          | ≥0.15 µg/ml** | 98.4%                 | 100%                             | -1.56 (-2.41, -0.71)                 |
|          | ≥1.00 µg/ml** | 84.7%                 | 88.9%                            | -4.15 (-6.93, -1.38)                 |

\*Mills, Elaine, et.al., *Vaccine*, 1998, Vol. 16, No. 6

\*\* Data on file